

Hemorrhage Control




COMBAT MEDIC ADVANCED SKILLS TRAINING (CMAST)

Introduction


- Review methods of hemorrhage control in a tactical environment.
- Hemorrhage is the leading cause of preventable death on the battlefield.
- Hemorrhage control save lives.
- New Hemostatic agents available.

Cardiovascular System



Blood Vessels

- Arteries
- Arterioles
- Capillaries
- Venules
- Veins




Pulses

- Left ventricle contracts.
- Peripheral pulses:
 - Radial
 - Brachial
 - Posterior tibial
 - Dorsalis pedis
- Central pulses:
 - Carotid
 - Femoral



Blood

- Adult body:
 - Contains approximately 5 to 6 liters of blood
 - Loss of 1 pint of blood without harmful effects
 - Loss of 2 pints may cause shock
- Three phases of hemostasis:
 - Vascular spasm
 - Platelet plug formation
 - Blood clotting (coagulation cascade)



Hemorrhage

- Pulse vs. Blood Pressure.
- How long until there are changes?
- Young healthy adults compensate for long periods, then decompensate rapidly.
- At what blood pressure do casualties lose consciousness?
 - @ 50 mm Hg

Clinical Signs of Acute Hemorrhage

Class	% Blood Loss	Clinical Signs
I	Up to 750 ml (15%)	Slight increase in HR; no change in BP or respirations
II	750-1500 ml (15-30%)	Increased HR and respirations; increased diastolic BP; anxiety, fright or hostility
III	1500-2000 ml (30-40%)	Increased HR and respirations; fall in systolic BP; significant AMS
IV	>2000 (>40%)	Severe tachycardia; severe lowering of BP; cold, pale skin; severe AMS

Sources of Hemorrhage

- External:
 - Visible blood is hard to estimate
- Internal:
 - May be hidden within the torso or even in the extremities secondary to fractures

Sources of External Bleeding

- Arterial:
 - Rapid, profuse and pulsating
 - Bright red in color
- Venous:
 - Steady flow
 - Dark red or maroon in color
- Capillary:
 - Slow and oozing
 - Often clots spontaneously



Extremity Hemorrhage



Click on picture for video
CMAST

Internal Signs of Hemorrhage

- Soft tissue bruising.
- Abdominal tenderness.
- Hemoptysis.
- Hematemesis.
- Melena.



Injured Internal Organs



Hemorrhage Control

- Assess the tactical situation.
- Expose the wound.
- Attempt to control the bleeding with direct pressure or a pressure dressing.



THE EMERGENCY BANDAGE




Trauma / Wound Compression Dressing

The Emergency Bandage is the first major innovation of its kind in over 50 years!

- NSN # 6510-01-460-0849 - FCP-01 - 4"
- NSN # 6510-01-492-2275 - FCP-02 - 6"
- NSN # 6510-01-515-7528 - FCP-05 - 6" +mobile pad

Hemorrhage Control



Click in box for video

CMAST

Hemorrhage Control

- Life-threatening arterial bleeding (amputation) may require early use of a tourniquet.
- If under enemy fire or in a dangerous position rapidly apply a tourniquet and move casualty to cover.

Tourniquets

- Several new tourniquets have been selected as primary means to control hemorrhage in combat.



Combat Application Tourniquet

- The C-A-T was selected as the primary tourniquet for every soldier.



SELF ADHERING BAND

WINDLAS
S

WINDLASS STRAP

C-A-T Step 1



Place the wounded extremity through the loop of the Self-adhering Band

C-A-T Step 2



**Place tourniquet
above the injury
site**

C-A-T Step 3

Pull the free-running end of the Self-adhering Band tight and securely fasten it back on itself.



C-A-T Step 4

Adhere Self-adhering Band completely around the limb until the clip is reached.



C-A-T Step 5

**Twist the
Windlass Rod
until the
bleeding has
stopped.**



C-A-T Step 6

Lock the Rod in place with the Windlass Clip



C-A-T Step 7



For small extremities, continue to adhere the Self-adhering Band around the extremity and over the Windlass Rod.

C-A-T Step 8

Grasp the Windlass Strap, pull it tight, and adhere it to the velcro on the Windlass Clip.



C-A-T Tourniquet



The C-A-T Tourniquet is now ready for transport.

C-A-T Tourniquet

NOTE:

The friction adaptor buckle is not necessary for proper C-A-T application to an arm. It **MUST** be used with two hands when applying to a leg.



C-A-T: Lower Extremity

To use, wrap the Self-adhering Band through the friction adaptor buckle.



C-A-T: Lower Extremity

This prevents the Self-adhering Band from loosening during transport.



C-A-T Tourniquet

C-A-Tourniquet
Arm Application

Click in box for video
CMAST

C-A-T: Lower Extremity



C-A-T technique
leg application

Click in box for video
CMAST

SOFTT



SOFTT Application

- Similar to the CAT:
 - Slide loop over extremity
 - Pull strap tight
 - Twist windlass until bleeding stops
 - Latch the windlass with one of the tri-rings
 - Tighten the safety screw



SOFTT Application



One-handed application




Two-handed application

Improvised Tourniquet

- Place cravat between heart and wound.
- Tie a half-knot on upper surface.
- Place a short stick on half-knot.
- Tie a square knot on top of stick.
- Twist stick (windlass) to tighten.
- UNTIL BLEEDING STOPS.
- Secure windlass to prevent unwinding.



Improvised Tourniquet



Tourniquet Principles

- Never cover a tourniquet.
- Mark a “T” on the casualty's forehead or somewhere obvious (sharpie pen).
- In combat when the tactical situation allows, loosening a tourniquet is appropriate.

Tourniquet Removal

- Once the tactical situation allows, tourniquets should be loosened and other methods to stop bleeding applied.
 - Direct pressure - pressure dressing
 - HemCon Chitosan Bandage
 - QuikClot powder

Tourniquet Removal

- When loosening a tourniquet, do not remove it from the limb.
- If the tourniquet has been in place for > 6 hours, do not remove.
- If fluid resuscitation is required, it should be accomplished before the tourniquet is removed.
- Tourniquets are very painful, provide pain medications as needed.

Tourniquet Removal

- If tourniquet has been in place for only 1-2 hours, loosening and using other methods to control hemorrhage can salvage limbs.
- Remember: if unable to control hemorrhage by other means, re-tighten the tourniquet.
- It is better to sacrifice the limb than to lose a life to hemorrhage.

Amputation

- Apply a pressure dressing to cover the end of the stump.
- Kerlix and 6" Ace wrap for effective pressure dressing.
- Rinse amputated part free of debris.
- Wrap loosely in saline-moistened sterile gauze.

Preservation of Amputation Parts

- Seal amputated part in a plastic bag or cravat.
- Place in a cool container; do not allow to freeze.
- Never place an amputated part in water.
- Never place amputated part directly on ice.
- Never use dry ice to cool an amputated part.

Hemostatic Agents

- HemCon® Chitosan Bandage.
- QuikClot® Hemostatic Powder.

Chitosan Hemostatic Dressing



- Hold the foil over-pouch so that instructions can be read. Identify unsealed edges at the top of the over-pouch.

Chitosan Hemostatic Dressing



- Peel open over-pouch by pulling the unsealed edges apart.

Chitosan Hemostatic Dressing



- Trap dressing between bottom foil and non-absorbable green/black polyester backing with your hand and thumb.

Chitosan Hemostatic Dressing



- Hold dressing by the non-absorbable polyester backing and discard the foil over-pouch. Hands must be dry to prevent dressing from sticking to them.

Chitosan Hemostatic Dressing



Chitosan Hemostatic Dressing

- Place the light colored sponge portion of the dressing directly to the wound area with the most severe bleeding. Apply pressure for 2 minutes or until the dressing adheres and bleeding stops. Once applied and in contact with the blood and other fluids, the dressing cannot be repositioned.
- A new dressing should be applied to other exposed bleeding sites; each new dressing must be in contact with tissue where bleeding is heaviest. Care must be taken to avoid contact with the casualty's eyes.


Chitosan Hemostatic Dressing

- If dressing is not effective in stopping bleeding after 4 minutes, remove the original and apply a new dressing. Additional dressings cannot be applied over ineffective dressings.
- Apply a battle dressing/bandage to secure a hemostatic dressing in place.
- Hemostatic dressings should only be removed by responsible persons after evacuation to the next level of care.



Chitosan HCD Porcine Abdominal Aorta Punch

Click on picture for video
CMAST





CMAST

55

QuikClot

- Warning: Avoid contact with wet skin; product reacts with small amounts of water and can cause burning.
- Stop burning by brushing away granules and flooding area with large volume of water.
- If ingested, immediately drink two or more glasses of water.

QuikClot

- Directions:

- 1-Apply direct firm pressure to wound using sterile dressing or best available substitute
- 2-If bleeding is stopped or nearly stopped after approximately 1 minute of pressure, wrap and tie bandage to maintain pressure on wound
- 3-If moderate to severe bleeding continues, hold pack away from face and tear open at tabs


QuikClot

- 4-Use wiping motion to remove gauze and excess blood – immediately start a slow pouring of one QuikClot packet directly onto the wound. Stop pouring as soon as dry granules cover the wound area
- 5-Use only enough QuikClot to stop bleeding. If bleeding continues open a second packet of QuikClot and continue to use as directed

QuikClot

6-Reapply firm pressure to QuikClot covered wound using sterile gauze. Wrap and tie bandage to maintain pressure

QuikClot



Click in box for video
CMAST

Treatment Goals

- Hemorrhage control continues to be the priority in battlefield care.
- Hemorrhage is the leading cause of preventable death on the battlefield.
- Our focus must be on stopping soldiers from bleeding to death on the battlefield.

Questions?

